

Effective Lagrangian with valence gluons in extended $SU(2)$ QCD model

Y. M. Cho^{1,2}, Haewon Lee³, D. G. Pak^{1,4}

¹⁾Asia Pacific Center for Theoretical Physics, Seoul 130-012, Korea

²⁾Department of Physics, College of Natural Sciences, Seoul National University, Seoul 151-742, Korea

³⁾Department of Physics, Chungbuk National University, Cheongju, Chungbuk 361-763, Korea

⁴⁾Department of Theoretical Physics, Uzbekistan National University, Tashkent 700-174, Uzbekistan

ymcho@yongmin.snu.ac.kr, hwlee@phys.chungbuk.ac.kr, dmipak@mail.apctp.org

The effective action with homogeneous valence (off-diagonal) gluons as background fields in the extended $SU(2)$ model of QCD is obtained in one-loop approximation. We keep the manifest gauge and Lorentz invariance during whole calculation by using a special gauge-fixing recipe and taking into account the quartic interaction terms in the original Lagrangian as well. It has been shown that the effective Lagrangian gains a positive imaginary part.

PACS numbers: 12.38.-t, 11.15.-q, 12.38.Aw, 11.10.Lm

The establishment of color confinement and derivation of the low energy effective theory from the first principles is one of important problems in quantum chromodynamics (QCD). For last years the idea of monopole condensation which can provide the color confinement mechanism [1,2] has been supported in analytical studies [3] and lattice simulations [4]. Recently it was shown [5] in the framework of the extended $SU(2)$ QCD model [6] that the effective theory obtained after integrating out all dynamical degrees of freedom can describe the confinement phase with the monopole condensation and dynamical gauge symmetry breaking. The later provides the effective mass generation and gives the origin for the kinetic term in the Skyrme-Faddeev effective Lagrangian [7,8,5]. It was proposed [5] that the effective action has a new duality symmetry which is closely related with the vacuum stability in the case of a pure chromomagnetic background whereas in the presence of a chromoelectric background the corresponding vacuum becomes unstable due to the imaginary part of the effective Lagrangian. The imaginary part for chromomagnetic and chromoelectric background in standard $SU(2)$ model of QCD was considered in [12–15].

In the present paper we study further the quantum properties of the extended $SU(2)$ model of QCD and calculate the one-loop effective action with covariant constant valence (off-diagonal) gluons as background fields. Notice, that the extended $SU(2)$ model of QCD [6] incorporates naturally the topological scalar field which insures the off-diagonal gluon to be represented as a gauge covariant object in the extended gauge theory. This allow us to treat the effective action with off-diagonal gluons in a consistent gauge invariant manner. Our effective Lagrangian with covariant constant valence gluons reveals a non-trivial vacuum but possesses a positive imaginary part. We maintain the explicit Lorentz and gauge invariance during calculation of the effective Lagrangian

and take into account the contribution which is coming from the quartic gluon interaction terms in the classical Lagrangian.

We will follow the original papers [6] and remind first the main lines of the extended $SU(2)$ model of QCD. The principal construction in that model is a generalized parametrization for the $SU(2)$ gauge connection implemented with a scalar field $\hat{n}$ which describes the topological degrees of freedom of the non-Abelian gauge theory

$$\begin{aligned}\vec{A}_\mu &= A_\mu \hat{n} - \frac{1}{g} \hat{n} \times \partial_\mu \hat{n} + \vec{X}_\mu \\ &= \hat{A}_\mu + \vec{X}_\mu, \quad (\hat{n}^2 = 1).\end{aligned}\quad (1)$$

The restricted connection $\hat{A}_\mu$ still represents an $SU(2)$ connection which undergoes the full $SU(2)$ group gauge transformations and at the same time maintains all topological characteristics of the original unconstrained non-Abelian potential. The isolated singularities of $\hat{n}$ define the second homotopy group $\pi_2(S^2)$ which describes the non-Abelian monopoles from the different topological sectors. Indeed, $\hat{A}_\mu$ with $A_\mu = 0$ and $\hat{n} = \hat{r}$ leads to Wu-Yang point-like monopole solution [9,10]. Besides that, with the S^3 compactification of the three-dimensional space R^3 , the all mappings $\hat{n}$ are characterized by the Hopf invariant $\pi_3(S^2) \simeq \pi_3(S^3)$ which classifies the topologically distinct vacua in the non-Abelian gauge theory [11,2]. Notice, that since the $SU(2)$ connection space is an affine space one can extend the restricted connection $\hat{A}_\mu$ by adding a general covariant vector $\vec{X}_\mu$ (so-called “valence gluons” [6]) which corresponds to the off-diagonal components of the gauge potential in the standard $SU(2)$ QCD.

Under the infinitesimal $SU(2)$ gauge transformation

$$\delta \hat{n} = -\vec{\theta} \times \hat{n}, \quad \delta \vec{A}_\mu = \frac{1}{g} \vec{D}_\mu \vec{\theta}, \quad (2)$$

one has the following transformation rules for the Abelian component A_μ (“photon”), the restricted connection $\hat{A}_\mu$ and valence gluons $\vec{X}_\mu$

$$\begin{aligned}\delta A_\mu &= \frac{1}{g} \hat{n} \cdot \partial_\mu \vec{\theta}, & \delta \hat{A}_\mu &= \frac{1}{g} \hat{D}_\mu \vec{\theta}, \\ \delta \vec{X}_\mu &= -\vec{\theta} \times \vec{X}_\mu,\end{aligned}\quad (3)$$

here

$$\hat{D}_\mu = \partial_\mu + g \hat{A}_\mu \quad (4)$$

is a covariant derivative of the restricted theory. It should be noted that the restricted connection $\hat{A}_\mu$ itself keeps the whole $SU(2)$ gauge invariance providing by this the gauge independent structure of the Abelian projected effective theory. The Lagrangian of the extended Yang-Mills theory can be written as follows [6]

$$\begin{aligned}\mathcal{L}_0 &= -\frac{1}{4} \vec{F}_{\mu\nu}^2 = -\frac{1}{4} \hat{F}_{\mu\nu}^2 - \frac{1}{2} \hat{F}_{\mu\nu} \cdot \hat{n} X_{\mu\nu} \\ &\quad - \frac{1}{4} X_{\mu\nu}^2 - \frac{1}{4} (\hat{D}_\mu \vec{X}_\nu - \hat{D}_\nu \vec{X}_\mu)^2,\end{aligned}\quad (5)$$

where the main field strengths for the “photon” A_μ , magnetic potential $\vec{C}_\mu$ and valence gluons $\vec{X}_\mu$ are defined by

$$\begin{aligned}\vec{F}_{\mu\nu} &= \hat{F}_{\mu\nu} \hat{n} + \hat{D}_\mu \vec{X}_\nu - \hat{D}_\nu \vec{X}_\mu + g \vec{X}_\mu \times \vec{X}_\nu, \\ \hat{F}_{\mu\nu} &= (F_{\mu\nu} + H_{\mu\nu}) \hat{n}, \\ F_{\mu\nu} &= \partial_\mu A_\nu - \partial_\nu A_\mu, \\ X_{\mu\nu} &= g \hat{n} \cdot (\vec{X}_\mu \times \vec{X}_\nu), \\ H_{\mu\nu} &= -\frac{1}{g} \hat{n} \cdot (\partial_\mu \hat{n} \times \partial_\nu \hat{n}) \\ &= (\partial_\mu \vec{C}_\nu - \partial_\nu \vec{C}_\mu + g \vec{C}_\mu \times \vec{C}_\nu) \cdot \hat{n}.\end{aligned}\quad (6)$$

The vector connection $\vec{C}_\mu \equiv -\frac{1}{g} \hat{n} \times \partial_\mu \hat{n}$ is exactly the magnetic potential which corresponds to the classical Wu-Yang monopole solution [9] or to Dirac string after making gauge transformation to a singular magnetic gauge [6]. In the magnetic gauge the magnetic potential for the monopole solution is represented explicitly by an Abelian part of the full connection. To pass to the magnetic gauge it is more convenient to decompose the vector fields $\vec{C}_\mu$ and $\vec{X}_\mu$ in the orthonormal basis $(\hat{n}^a, n^3 \equiv \hat{n}; a = 1, 2)$ instead of explicit rotating the internal space of $SU(2)$ as it was defined in [6]

$$\vec{C}_\mu = C_\mu^a \hat{n}^a, \quad \vec{X}_\mu = \vec{X}_\mu^a \hat{n}^a. \quad (7)$$

After introducing complex notations for the vector fields $X_\mu = \frac{1}{\sqrt{2}}(X_{1\mu} + iX_{2\mu})$, $\bar{X}_\mu = (X_\mu)^*$ one can rewrite the classical Lagrangian $\mathcal{L}_0$ as

$$\begin{aligned}\mathcal{L}_0 &= -\frac{1}{4}(F_{\mu\nu} + H_{\mu\nu})^2 - \frac{1}{2}|D_\mu X_\nu - D_\nu X_\mu|^2 \\ &\quad + ig(F_{\mu\nu} + H_{\mu\nu})\bar{X}_\mu X_\nu - V(X^4), \\ V(X^4) &= \frac{g^2}{2}[(\bar{X}X)^2 - \bar{X}^2 X^2], \\ D_\mu &= \partial_\mu + ig(A_\mu + C_\mu), \\ C_\mu &= -\frac{1}{g}\hat{n}_1\partial_\mu\hat{n}_2.\end{aligned}\quad (8)$$

The magnetic field strength $H_{\mu\nu}$ can be rewritten in the magnetic gauge as follows

$$H_{\mu\nu} = \partial_\mu C_\nu - \partial_\nu C_\mu. \quad (9)$$

We have two types of gauge transformations under which the original Lagrangian is invariant:

(I) the active type transformations defined by Eqs. (2) and (3) and (II) the passive ones which have the following form

$$\begin{aligned}\delta C_\mu &= 0, \\ \delta A_\mu &= \frac{1}{g} \partial_\mu \theta_3 + i\bar{\theta} X_\mu - i\theta \bar{X}_\mu, \\ \delta X_\mu &= \frac{1}{g} D_\mu \theta - i\theta^3 X_\mu.\end{aligned}\quad (10)$$

It should be noted that the passive type gauge transformations possess a structure of the complexified $U(1)$ gauge group and in addition we have also the Abelian dual $U_m(1)$ magnetic gauge symmetry for the magnetic potential C_μ [6].

Let us consider the generating functional for connected Green functions with the sources corresponding to the Abelian and off-diagonal gluons

$$\begin{aligned}W[J_\mu, \vec{J}_\mu] &= \int DA_\mu D\vec{X}_\mu \exp[-i \int (\frac{1}{4} \vec{F}_{\mu\nu}^2 \\ &\quad + A_\mu J_\mu + \vec{X}_\mu \cdot \vec{J}_\mu) d^4x].\end{aligned}\quad (11)$$

We split the gauge fields into the classical background $A_{0\mu}, X_{0\mu}$ and quantum parts A'_μ, X'_μ

$$\begin{aligned}A_\mu &= A_{0\mu} + A'_\mu, \\ X_\mu &= X_{0\mu} + X'_\mu.\end{aligned}\quad (12)$$

The magnetic potential C_μ (or equivalently the topological scalar field $\hat{n}$) is treated as a classical object since it does not contain dynamical degrees of freedom in the original theory. Notice that formally we can include the magnetic potential C_μ into the classical part $A_{0\mu}$ of the Abelian gauge component since the C_μ enters the covariant derivatives $D_\mu, \bar{D}_\mu$ in additive manner everywhere. In computation of functional determinants in the non-Abelian gauge theory while keeping the manifest Lorentz

invariance one encounters an obstacle of calculating the matrix determinants with respect to Lorentzian and internal group indices. To overcome this technical difficulty we choose the special gauge-fixing functions f, h

$$\begin{aligned} f &= D_{0\mu}X'_\mu - igA'_\mu X_{0\mu}, \\ h &= \partial_\mu A'_\mu + igX_{0\mu}\bar{X}'_\mu - ig\bar{X}_{0\mu}X'_\mu, \end{aligned} \quad (13)$$

where $D_{0\mu} = \partial_\mu + ig(A_{0\mu} + C_\mu)$ is a $U(1)$ covariant derivative with background Abelian fields. This choice allows us to factorize the functional determinants in Lorentzian indices. Using the passive type gauge transformations and applying the 't Hooft trick of inserting the unit into the generating functional we obtain the gauge-fixing Lagrangian (in Feynman gauge)

$$\mathcal{L}_{gf} = -f f^* - \frac{1}{2}h^2 \quad (14)$$

and the Faddeev-Popov ghost determinant $\text{Det}M_{FP}$ in a standard way. We calculate the effective action in one-loop approximation and consider the background valence gluons which are covariant constant under the transformations of the stability subgroup $U(1)$

$$D_{0\mu}X_\nu = 0. \quad (15)$$

Since the background Abelian field $A_{0\mu} + C_\mu$ appears everywhere only in terms of the covariant background derivative $D_{0\mu}$ and background field strength $F_{0\mu\nu}$ we can rewrite the effective Lagrangian in an explicit covariant form. Our choice of the gauge-fixing functions, Eq. (13), simplifies crucially the kinetic term structure

$$\begin{aligned} L_0 + L_{gf} &= \frac{1}{2}A'_\mu[g_{\mu\nu}(\square - 2a)]A'_\nu \\ &\quad + 2igA'_\mu(X_{0\nu}\bar{D}_{0\nu}\bar{X}'_\mu - \bar{X}_{0\nu}D_{0\nu}X'_\mu) \\ &\quad + \bar{X}'_\mu[g_{\mu\nu}(D_0D_0 - a) + 2igF_{0\mu\nu}]X'_\nu \\ &\quad + 2g^2\bar{X}'_\mu(\bar{X}_{0\mu}X_{0\nu} - X_{0\mu}\bar{X}_{0\nu})X'_\nu \\ &\quad + \frac{g^2}{2}\bar{X}'^2X_0^2 + \frac{g^2}{2}X'^2\bar{X}_0^2, \end{aligned} \quad (16)$$

where $a = g^2\bar{X}_{0\mu}X_{0\mu}$ and we keep only terms quadratic in quantum fields which are enough at one-loop level. The Faddeev-Popov matrix operator is

$$\begin{aligned} M_{FP} &= \frac{\delta(f, \bar{f}, h)}{\delta(\theta, \bar{\theta}, \theta^3)} \\ &= \frac{1}{g} \begin{pmatrix} D_0D_0 - a & g^2X_0^2 & -2igX_{0\mu}\partial_\mu \\ g^2\bar{X}_0^2 & \bar{D}_0\bar{D}_0 - a & 2ig\bar{X}_{0\mu}\partial_\mu \\ -2ig\bar{X}_{0\mu}D_{0\mu} & 2igX_{0\mu}\bar{D}_{0\mu} & \square - 2a \end{pmatrix}. \end{aligned} \quad (17)$$

We consider the case of $U(1)$ covariant constant valence gluon background. The integration over Abelian field A_μ results in a functional determinant

$$\text{Det}^{-\frac{1}{2}}K_A = \text{Det}^{-\frac{1}{2}}[g_{\mu\nu}(\square - 2a)] \quad (18)$$

and additional non-local terms which will contribute to the effective action after subsequent integrating out the quantum parts X'_μ of the valence gluons. The final expression for the functional determinant obtained after performing the integration over Abelian and off-diagonal gauge fields is the following

$$\begin{aligned} \text{Det}^{-\frac{1}{2}}K_{\mu\nu} &= \text{Det}^{-\frac{1}{2}} \begin{pmatrix} \tilde{A} & \tilde{B} \\ \tilde{C} & \tilde{D} \end{pmatrix}, \\ \tilde{A}_{\mu\nu} &= g_{\mu\nu}(D_0D_0 - a + \frac{\alpha}{\square - 2a}) + 2ig(F_{0\mu\nu} + X_{0\mu\nu}), \\ \tilde{B}_{\mu\nu} &= g_{\mu\nu}(g^2X_0^2 - \frac{\beta}{\square - 2a}) \equiv g_{\mu\nu}B, \\ \tilde{C}_{\mu\nu} &= (\tilde{B}^*)_{\mu\nu} \equiv g_{\mu\nu}C, \quad \tilde{D}_{\mu\nu} = \tilde{A}_{\mu\nu}^*. \end{aligned} \quad (19)$$

where

$$\begin{aligned} \alpha &= 4g^2\bar{X}_{0\mu}X_{0\nu}D_{0\mu}D_{0\nu}, \\ \beta &= 4g^2X_{0\mu}X_{0\nu}\bar{D}_{0\mu}\bar{D}_{0\nu}. \end{aligned} \quad (20)$$

So that the one-loop contribution to the effective action is given by

$$\Delta S_{eff} = -\frac{1}{2}\text{Tr} \ln K_A - \frac{1}{2}\text{Tr} \ln K_{\mu\nu} + \text{Tr} \ln M_{FP}. \quad (21)$$

For simplicity we consider the effective Lagrangian with off-diagonal background gauge fields only, so from now on we set $A_{0\mu} = 0$. The determinant of the matrix $K_{\mu\nu}$ can be simplified to the next form

$$\begin{aligned} \text{Det}K_{\mu\nu} &= \text{Det}(\tilde{A}\tilde{D} - \tilde{B}\tilde{C}) \\ &= \text{Det}[g_{\mu\nu}(A^2 - BC) + 4X_{0\mu\rho}X_{0\rho\nu}], \end{aligned} \quad (22)$$

where

$$A = \square - a + \frac{\alpha}{\square - 2a}. \quad (23)$$

The field strength $X_{\mu\nu}$ has a useful cyclic property

$$\begin{aligned} X_{\mu\nu}X_{\nu\rho}X_{\rho\sigma} &= -\frac{1}{2}\chi^2X_{\mu\sigma}, \\ \chi &= \sqrt{X_{0\mu\nu}X_{0\mu\nu}}. \end{aligned} \quad (24)$$

Using this relation one can calculate the part of the determinant in Eq. (22) with respect to Lorentzian indices. After some simplifications one obtains

$$\begin{aligned} \ln \text{Det}K_{\mu\nu} &= 2\text{tr} \ln(A^2 - BC) + 2\text{tr} \ln(A^2 - BC - 2\chi^2), \end{aligned} \quad (25)$$

where “tr” involves only the remaining functional trace.

The Faddeev-Popov determinant is simplified in a similar manner

$$\text{Det} M_{FP} = \text{Det}(A^2 - BC) \text{Det}(\square - 2a). \quad (26)$$

Summing up contributions from all determinants one obtains the simple expression for the one-loop contribution to the effective action

$$\Delta S_{eff} = -\text{tr} \ln(A^2 - BC - 2\chi^2) - \text{tr} \ln(\square - 2a). \quad (27)$$

This expression has no neither Lorentzian nor internal group indices and is quite suitable for calculation of the remaining functional traces. Passing to the Euclidian space-time and making Fourier transformation we can fulfill the integration over momentum with using a cut-off regularization scheme. To obtain the final expression for the effective Lagrangian one needs to know only the functional dependence on the valence gluons. The easiest way to find the functional form of the effective Lagrangian is to replace the complex vectors $X_\mu, \bar{X}_\mu$ with a pair of real vectors in Euclidian space-time

$$X_{\mu E} \rightarrow V_\mu, \quad \bar{X}_{\mu E} \rightarrow W_\mu. \quad (28)$$

In four dimensional Euclidian spherical coordinates the momentum k_μ has the following projections

$$k_\mu = \begin{pmatrix} k \cos \theta_1 \\ k \sin \theta_1 \cos \theta_2 \\ k \sin \theta_1 \sin \theta_2 \cos \phi \\ k \sin \theta_1 \sin \theta_2 \sin \phi \end{pmatrix}. \quad (29)$$

One can choose the coordinate system in the 4-dimensional Euclidean space-time in such a manner that the vectors V_μ and W_μ will have the next components

$$V_\mu = (V, 0, 0, 0), \quad W_\mu = (W \cos \psi, W \sin \psi, 0, 0), \quad (30)$$

here, we introduced the angle ψ between vectors V_μ, W_μ . The knowledge of the angle dependence allows us to restore the scalar product terms $(\bar{X}X)$ from the corresponding terms $VW \cos \psi$ in the final expression after completing whole calculation. It is possible to proceed through all calculations with the original real vectors $X_{1\mu}, X_{2\mu}$ as well. But for our purpose to calculate the effective Lagrangian with a constant background the replacement defined by Eq. (28) is more convenient. With these notations one can write down the one-loop contribution to the effective action as follows

$$\begin{aligned} \Delta S_{eff} = & -\text{tr} \ln[(k^2 - a)^2(k^2 - 2a) + 2\alpha(k^2 - a) \\ & - (4a^2 - 3g^4 V^2 W^2)(k^2 - 2a) \\ & - g^2 W^2 \beta - g^2 V^2 \bar{\beta}], \end{aligned} \quad (31)$$

where

$$\begin{aligned} a &= -g^2 q VW, \\ q &= \cos \psi, \\ \alpha &= -4g^2 VW k^2 \cos \theta_1 (q \cos \theta_1 + \sin \psi \sin \theta_1 \cos \theta_2), \\ \beta &= -4g^2 V^2 k^2 \cos^2 \theta_1, \\ \bar{\beta} &= -4g^2 W^2 k^2 (q \cos \theta_1 + \sin \psi \sin \theta_1 \cos \theta_2)^2. \end{aligned} \quad (32)$$

One can calculate explicitly the leading logarithmic part in the one-loop effective Lagrangian using a cut-off regularization. After performing the integration over $k, \theta_1, \theta_2, \phi$ one results in

$$\begin{aligned} L_{eff} = & -\frac{g^2}{4} \chi^2 - g^2 \Lambda^2 (\bar{X}_0 X_0) \\ & - \frac{11g^4}{96\pi^2} \chi^2 (\ln[\frac{g^2 \chi}{\Lambda^2}] - c) + \frac{11ig^4}{192\pi} \chi^2, \end{aligned} \quad (33)$$

where c is a number parameter, and we omitted the quartic terms proportional to Λ^4 . To calculate the number factor c and the imaginary part of the effective Lagrangian we used numerical computation with considering an analytical continuation in the parameter q from $q = 1$ to $q = 0$ which corresponds to analytical continuation of the χ^2 to negative values. Since the final expression possesses a covariant form one can calculate the number parameter $c = 0.3769...$ and, what is more important, the imaginary part of the effective Lagrangian. After subtracting infinities within the minimal subtraction scheme and proper mass and charge renormalization we obtain a final expression for the one-loop renormalized effective Lagrangian with valence gluons

$$\begin{aligned} \mathcal{L}_{eff}^{ren} = & -\frac{g^2}{4} \chi^2 - m_R^2 (\bar{X}_0 X_0) \\ & - \frac{11g^4}{96\pi^2} \chi^2 (\ln[\frac{g^2 \chi}{\mu^2}] - \frac{1}{2}) + \frac{11ig^4}{192\pi} \chi^2. \end{aligned} \quad (34)$$

It should be stressed that the sign of the imaginary part is fixed by the causal structure which is under the control by putting the infinitesimal quantity $-i\epsilon$ in propagators. For the case $q = 0$ one has checked that the real part of the function under the logarithm in Eq. (31) is negative for all admissible values of k_μ . This determines uniquely the sign of the imaginary part in the effective Lagrangian. We can keep also the renormalised mass parameter m_R since in the presence of the magnetic potential C_μ the off-diagonal gluons behave like a charged matter and the mass term does not spoil the gauge invariance. In the case of $m_R = 0$ the vacuum energy density has a non-trivial vacuum at the non-zero value for the classical field strength χ

$$\langle \chi \rangle = \frac{\mu^2}{g^2} \exp(-\frac{24\pi^2}{11g^2}). \quad (35)$$

This minimum does not correspond to a stable vacuum due to the presence of the imaginary part in the effective Lagrangian.

Notice, that our final expression from the effective Lagrangian, Eq. (34), has a manifest gauge and Lorentz invariant form and the number factor in front of the logarithm leads to a correct negative β -function of QCD in agreement with the renormalization group equation approach.

-
- [1] Y. Nambu, Phys. Rev. **D10**, 4262 (1974); S. Mandelstam, Phys. Rep. **23C**, 245 (1976); A. Polyakov, Nucl. Phys. **B120**, 429 (1977).
 - [2] G. 't Hooft, Nucl. Phys. **B190**, 455 (1981).
 - [3] Z. Ezawa, and A. Iwazaki, Phys. Rev. **D25**, 2681 (1982); T. Suzuki, Prog. Theor. Phys. **80**, 929 (1988); H. Suganuma, S. Sasaki, and H. Toki, Nucl. Phys. **B435**, 207 (1995); K.-I. Kondo, Phys. Rev. **D57**, 7467 (1998); **D58**, 105016 (1998).
 - [4] A. Kronfeld, G. Schierholz, and U. Wiese, Nucl. Phys. **B293**, 461 (1987); T. Suzuki, and I. Yotsuyanagi, Phys. Rev. **D42**, 4257 (1990); J. Stack, S. Neiman, and R. Wensley, Phys. Rev. **D50**, 3399 (1994); H. Shiba, and T. Suzuki, Phys. Lett. **B333**, 461 (1994); G. Bali, V. Bornyakov, M. Müller-Preussker, and K. Schilling, Phys. Rev. **D54**, 2863 (1996).
 - [5] Y. M. Cho, and D. G. Pak, *Proc. of TMU-YALE Symposium on Dynamics of Gauge Fields*, ed. by T. Appelquist, and H. Minakata (Univ. Acad. Press, Tokyo) 2000, 193-212, hep-th/0006051; J. Kor. Phys. Soc. **38**, 151 (2001).
 - [6] Y. M. Cho, Phys. Rev. **D21**, 1080 (1980); Phys. Rev. Lett. **46**, 302 (1981); Phys. Rev. **D23**, 2415 (1981).
 - [7] L. Faddeev, and A. Niemi, Nature **387**, 58 (1997); R. Battye, and P. Sutcliffe, Phys. Rev. Lett. **81**, 4798 (1998).
 - [8] E. Langmann, and A. Niemi, Phys. Lett. **B463**, 252 (1999); H. Gies, *Wilsonian effective action for SU(2) Yang-Mills theory with Cho-Faddeev-Niemi-Shabanov decomposition*, hep-th/0102026.
 - [9] T. T. Wu, and C. N. Yang, Phys. Rev. **D12**, 3845 (1975).
 - [10] Y. M. Cho, Phys. Rev. Lett. **44**, 1115 (1980); Phys. Lett. **B115**, 125 (1982).
 - [11] A. Belavin, A. Polyakov, A. Schwartz, and Y. Tyupkin, Phys. Lett. **59B**, 85 (1975); Y. M. Cho, Phys. Lett. **B81**, 25 (1979).
 - [12] G. K. Savvidy, Phys. Lett. **B71**, 133 (1977).
 - [13] N. K. Nielsen, and P. Olesen, Nucl. Phys. **B144**, 485 (1978);
 - [14] C. Rajiadaikos, Phys. Lett. **B100**, 471 (1981).
 - [15] V. Schanbacher, Phys. Rev. **D26**, 489 (1982).